LECTURE 2: 1-D PROBLEMS, LINEAR ANALYSIS, BIFURCATION, BI-FURCATION DIAGRAM

Contents

1	Intr	roduction	2
2	Line 2.1 2.2	earisation of 1-D systems Linearisation about fixed point x^* and linear stability analysis of x^*	2 2 4
3	Exa 3.1 3.2	Imple of linearisation and linear stability analysisLinearisation of the logistic equationLinear stability analysis of the fixed points	5 5 5
4	Existence and uniqueness of solutions of 1-D systems		
5	Impossibility of oscillations in 1-D systems		6
6	Bifurcation		7
	6.1	Saddle-node bifurcation	7
		6.1.1 Bifurcation diagram	8
		6.1.2 Not so clear example of a saddle-node bifurcation	8
		6.1.3 Normal form	10
	6.2	Transcritical bifurcation	10
	6.3	Pitchfork bifurcation	11
		6.3.1 Supercritical pitchfork bifurcation	12

1 Introduction



Figure 1: Phase portrait of a 1-D problem or a first-order problem. Fixed points are shown with the filled and empty bullets. The arrowheads show the direction of flow. The 1-D flow takes place on the *x*-axis.

We continue with the one-dimensional problems or the problems of flow on a line given in the form:

$$\dot{x} = f(x),\tag{1}$$

where function f(x) can be linear or nonlinear. Figure 1 shows the phase portrait of a 1-D problem. At times it is beneficial to simplify your problems in order to analyse them. This is especially true in case of problems that are described by nonlinear differential equations. Linearisation is a tool for analysing dynamics of nonlinear systems.

2 Linearisation of 1-D systems

2.1 Linearisation about fixed point x^* and linear stability analysis of x^*

In mathematics, <u>linearisation is finding a linear approximation to a function at a given point</u>. The linear approximation of a function is the first order Taylor expansion around the point of interest. In the study of dynamical systems, linearisation is a method for assessing the <u>local stability and type of a fixed point</u> of a system of nonlinear differential equations or discrete dynamical systems.

	SLIDES: 3, 4
Linearisation of 1-D systems	Linearisation of 1-D systems
The one-dimensional system is given by	Taylor series expansion about x^* of (4) results in
$\dot{x} = f(x). \tag{1}$	$\dot{\eta} = f(x) = f(x^* + \eta) \tag{5}$
The dynamics close to fixed point \boldsymbol{x}^* can be expressed as follows:	$= f(x^*) + \frac{f'(x^*)}{1!}(x^* + \eta - x^*) + \frac{f''(x^*)}{2!}(x^* + \eta - x^*)^2 + \dots $ (6)
$x(t) = x^* + \eta(t),$ (2)	$= f'(x^*)\eta + \frac{f''(x^*)}{2!}\eta^2 + \dots $ (7)
where $ \eta \ll 1$ is a small perturbation. The behaviour and change of solution x over time thus is	higher order terms, $O(\eta^2)$ $\approx f'(x^*)\eta.$ (8)
$\dot{x} = (x^* + \eta) = \dot{\eta}.$ (3)	If $f'(x^*) \neq 0$, then term $ f'(x^*)n \gg \left \frac{f''(x^*)}{\alpha}n^2\right $. Neglecting $O(n^2)$
At the same time (1) holds. This mans that the dynamics of small porturbations is the following:	yields the linearisation of the system about fixed point x^*
$\frac{1}{2} f(x) = f(x^* + x) $ (4)	$\dot{\eta} = s\eta,$ (9)
$\eta = f(x) = f(x^* + \eta). \tag{4}$	where $s=f^\prime(x^\ast)$ is simply the slope of function $f(x)$ evaluated at $x^\ast.$
D. Kartofelev YFX1560 3/9	D. Kartofelev YFX1560 4/9

Let's examine the dynamics of Eq. (1) close to its fixed point or points x^* . We assume the solution is in the following form:

$$x(t) = x^* + \eta(t), \tag{2}$$

where $|\eta| \ll 1$ is a small perturbation. The behaviour and change of solution x over time thus is

$$\dot{x} = (x^* + \eta) = \dot{\eta}. \tag{3}$$

At the same time it holds that $\dot{x} = f(x)$. The combination of these results gives

$$\dot{\eta} = f(x) = f(x^* + \eta) = \begin{bmatrix} \text{Taylor ser.} \\ \text{expansion} \\ \text{about } x^* \end{bmatrix} = f(x^*) + \frac{f'(x^*)}{1!}(x^* + \eta - x^*) + \frac{f''(x^*)}{2!}(x^* + \eta - x^*)^2 + \dots = \\ = \begin{bmatrix} f(x^*) = 0 \end{bmatrix} = f'(x^*)\eta + \underbrace{\frac{f''(x^*)}{2!}\eta^2 + \dots}_{\text{higher order terms, } O(\eta^2)} \approx f'(x^*)\eta. \quad (4)$$

If $f'(x^*) \neq 0$, then the term $|f'(x^*)\eta| \gg \left|\frac{f''(x^*)}{2!}\eta^2\right|$. Neglecting $O(\eta^2)$ yields the linearisation of the system about fixed point x^*

$$\dot{\eta} = s\eta,\tag{5}$$

where $s = f'(x^*)$ is simply the slope of function f(x) evaluated at x^* .

The linearised form (5) of the original system (1) is a familiar equation to us. Solution of Eq. (5) has the form

$$\eta(t) = \mathbf{e}^C \mathbf{e}^{st} = \eta_0 \, \mathbf{e}^{st},\tag{6}$$

where C is the constant of integration and η_0 is a suitable initial amplitude. The stability of this solution based on its behaviour is the following:

- If s > 0, then the solution is exponentially growing (exploding). We say that the solution dynamics and corresponding fixed point are **unstable**.
- If s < 0, then the solution is exponentially decaying. Asymptotically approaching a *stable* value $(\eta(t) \to 0 \text{ for } t \to \infty)$. We conclude that the solution dynamics and corresponding fixed point are **stable**.

This behaviour is also clearly seen on the example phase portrait shown in Fig. 2. The linearised phase portraits (shown in red) preserve the characteristics of the original flow (shown in black) in the close proximity of their respective fixed point x_1^* and x_2^* .



Figure 2: (Top) Phase portraits of the original flow described by Eq. (1) and its linearised counterparts given by Eq. (5). Linearisation is performed about fixed points x_1^* and x_2^* . The slopes s_1 and s_2 correspond to fixed points x_1^* and x_2^* , respectively. Linearised solutions are shifted to coincide with the positions of the fixed points. Linearised phase portraits are similar to the original flow only in close proximity to their respective fixed points. (Bottom) Aforementioned linearised solutions shown on their respective phase portraits.

If $s = f'(x^*) = 0$, then no information from linearisation can be obtained. The determination of stability type needs further analysis, i.e., additional terms in the Taylor series expansion (4) have to be considered.

2.2 Examples of cases where $f'(x^*) = 0$

Consider the following systems

$$\dot{x} = x^2,\tag{7}$$

$$\dot{x} = -x^2,\tag{8}$$

$$\dot{x} = x^3,\tag{9}$$

$$\dot{x} = -x^3. \tag{10}$$

Systems (7)–(10) share the same fixed point $x^* = 0$ and have different stability types, this can be clearly seen in Fig. 3. Systems (7) and (8) feature a new type of fixed point—the **half-stable fixed point**. The origin and nature of the half-stable fixed point will become clear below (Sec. 6.1).

Systems (9) and (10) feature **algebraic decay** $x(t) \sim t^{-\delta}$ near the fixed point (see Slide 5), as opposed to a more common (in the context of our course) **exponential decay** $x(t) \sim e^{-\gamma t}$, where γ is constant. We will revisit this idea in future lectures because algebraic decay may be a possible source of *confusion* in analysis of 2-D systems.



Figure 3: Phase portraits of Eqs. (7)-(10). Upper row of the graphs features a new type of fixed point—the **half-stable fixed point** that is stable on one side but not on the other, hence the half filled bullets.



Example of linearisation and linear stability analysis 3

3.1Linearisation of the logistic equation

The logistic equation has the form

$$\dot{x} = f(x) = rx\left(1 - \frac{x}{K}\right),\tag{11}$$

where $x \ge 0$ is the population size, r > 0 and K > 0 are the system parameters (see Lecture 1). Fixed points are the following:

$$\dot{x} = 0 \Rightarrow rx^* \left(1 - \frac{x^*}{K} \right) = 0 \Rightarrow \begin{cases} x_1^* = 0, \\ x_2^* = K. \end{cases}$$
(12)

In order to obtain linearisation in form (5), we calculate the slope of the right hand side of (11) and evaluate it at fixed points x_1^* and x_2^* . The slope is

$$f'(x) = \frac{\mathrm{d}}{\mathrm{d}x} \left[rx \left(1 - \frac{x}{K} \right) \right] = \frac{\mathrm{d}}{\mathrm{d}x} \left(rx - \frac{r}{K} x^2 \right) = r - \frac{2r}{K} x.$$
(13)

Evaluation of the obtained slope at x^* results in the following linearised system

$$\dot{\eta} = f'(x^*)\eta = \left(r - \frac{2r}{K}x\right)\Big|_{x=x^*}\eta.$$
(14)

In the case $x_1^* = 0$ we have

$$\dot{\eta} = \left(r - \frac{2r}{K}x\right)\Big|_{x=x_1^*=0} \eta = r\eta,$$
(15)

and in the case $x_2^* = K$ we have

$$\dot{\eta} = \left(r - \frac{2r}{K}x\right)\Big|_{x=x_2^*=K} \eta = (r - 2r)\eta = -r\eta.$$
(16)

The phase portrait shown in Fig. 4 shows the obtained linearised systems using the red graphs.



Figure 4: Stability of the fixed points is indicated by the hollow and filled bullets. Linearised results are shown with the red lines. $\dot{\eta} = r\eta$ for $x_1^* = 0$ and $\dot{\eta} = -r(\eta - K)$ for $x_2^* = K$, here the graph $\dot{\eta} = -r\eta$ (16) is shifted to coincide with the actual position of the fixed point $x_2^* = K$.

3.2Linear stability analysis of the fixed points

In order to evaluate the stability of the above fixed points we need to evaluate the signs of slopes $f'(x^*)$.

- For $x_1^* = 0$ we get: $f'(x^*) = \left(r \frac{2r}{K}x\right)\Big|_{x=x_1^*=0} = r$. The fixed point is **unstable** because the sign of
- the slope is **positive**. For $x_2^* = K$ we get: $f'(x^*) = \left(r \frac{2r}{K}x\right)\Big|_{x=x_2^*=K} = -r$. The fixed point is **stable** because the sign of the slope is **negative**.

Figure 4 shows the clear relationship between the slopes of the linearised results and the stability of the corresponding fixed points.

4 Existence and uniqueness of solutions of 1-D systems



backward time t.

5 Impossibility of oscillations in 1-D systems

The 1-D systems are limited in what they can describe. The possible behaviours of solution x(t) as $t \to \infty$, for $\dot{x} = f(x)$ are:

(i) $x(t) \to \pm \infty$ as $t \to \infty$.

(ii) $x(t) \to x^*$ as $t \to \infty$.

The behaviours that are not possible include: <u>oscillations</u>, <u>periodic</u> and <u>quasi-periodic solutions</u>, <u>chaos</u> (last two defined and explained in future lectures). Two mentioned examples of impossible behaviours are shown in Fig. 5



Figure 5: Examples of impossible behaviours in 1-D systems: (Left) oscillation with attenuation, (Right) harmonic oscillation.



Figure 6: Phase portrait of an arbitrary 1-D flow problem where $x_0 = x(0)$ is the initial condition.

Why is this true? Let's study Fig. 6 where the flow is always damped out at the stable fixed points for $t \to \infty$. By definition 1-D systems do not have inertial terms (\ddot{x}) this means that all forces are always balanced out by the damping, remember that \dot{x} term is proportional to losses or damping, see Eq. (1).

6 Bifurcation



Slide 7 gives the definition of a bifurcation. In this section our goal is to familiarise ourselves <u>with a selection</u> of basic bifurcation dynamics that occur in 1-D systems. They are:

- Saddle-node bifurcation
- Transcritical bifurcation
- Pitchfork bifurcation
 - Supercritical pitchfork bifurcation
 - Subcritical pitchfork bifurcation

6.1 Saddle-node bifurcation



Figure 7: Phase portraits of Eq. (17) undergoing a saddle-node bifurcation.

Basic mechanism for creation or destruction of fixed points. The *standard* example is

$$\dot{x} = r + x^2,\tag{17}$$

where r is the **control parameter**. Figure 7 shows phase portraits corresponding to different values of r. Fixed points exist only for $r \leq 0$ and they are given by

$$\dot{x} = 0 \quad \Rightarrow \quad r + x^{*2} = 0 \quad \Rightarrow \quad x^* = \pm i\sqrt{r}, \quad r \le 0, \tag{18}$$

where i is the imaginary unit. The **bifurcation point** value r = 0 and it occurs at spatial point x = 0, we write (x, r) = (0, 0). This example also shows the origin of the mysterious **half-stable fixed point** present in Eqs. (7) and (8). In those examples we simply caught the system at a moment of bifurcation.

A similar general behaviour, i.e., bifurcation dynamics, happens for the system with inverted parabola:

$$\dot{x} = r - x^2. \tag{19}$$

Here, the fixed points exists only for $r \ge 0$ their stability is the opposite compared to the previous example, and their values are the following:

$$\dot{x} = 0 \quad \Rightarrow \quad r - x^{*2} = 0 \quad \Rightarrow \quad x^* = \pm \sqrt{r}, \quad r \ge 0.$$

$$(20)$$

6.1.1 Bifurcation diagram

Bifurcation diagram is obtained by plotting the graph of fixed point x^* as a function of control parameter r, analytically expressed by (18). The bifurcation diagram of Eq. (17) is shown in Fig. 8. Compare Fig. 8 to the corresponding Fig. 7.



Figure 8: Bifurcation diagram of the saddle-node bifurcation. The unstable branch is shown with the dashed bold curve and the stable branch with the continuous bold curve.

The following numerical file shows the quantitatively accurate phase portraits and bifurcation diagrams discussed in this section.

NUMERICS: NB#1 Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (supercritical), pitchfork bifurcation (subcritical).

6.1.2 Not so clear example of a saddle-node bifurcation

Let's consider the following problem

$$\dot{x} = r + x - \ln(1 + x), \tag{21}$$

where r is the control parameter. It is impossible to determine the dynamics of the fixed points algebraically. Try it, and/or see the numerical file linked below.

NUMERICS: NB#1 Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (supercritical), pitchfork bifurcation (subcritical).

Note: The red error messages from Mathematica are left in intentionally. The symbolic calculation packages (CAS – computer algebra system) can not handle the problem using inverse functions.

We employ a graphical approach to find the fixed points, bifurcation point and their dependents on control parameter r. This means that we need to solve

$$\dot{x} = 0 \quad \Rightarrow \quad r + x - \ln(1 + x) = 0, \tag{22}$$

or equivalently

$$r + x = \ln(1 + x),$$
 (23)



Figure 9: (Left) Behaviour of functions (24) as the value of r is changed. (Right) Bifurcation happening for the tangent intersection of the line and the curve given by (24).

for variable x. We can use an arbitrary axis y to plot the left-hand side and the right-hand side of Eq. (23)

$$y = r + x$$
 and $y = \ln(1 + x)$. (24)

Figure 9 shows the behaviour of the line and the curve given by functions (24) as the value of r is changed. <u>A saddle-node bifurcation occurs</u> when the tangent intersection takes place. We write and it must hold

$$\begin{cases} r+x = \ln(1+x), \\ \frac{\mathrm{d}}{\mathrm{d}x}(r+x) = \frac{\mathrm{d}}{\mathrm{d}x}\ln(1+x). \end{cases}$$
(25)

As stated above the first equation in Sys. (25) is *hard* to solve algebraically, but <u>the second</u> one is easier. After taking the derivatives we get

$$1 = \frac{1}{1+x} \quad \Rightarrow \quad x^* = 0. \tag{26}$$

The corresponding r value can be found from the first equation in Sys. (25)

$$(r+x)|_{x=x^*=0} = \ln(1+x)|_{x=x^*=0},$$
(27)

$$r = \ln 1 = 0. \tag{28}$$

Thus, the bifurcation coordinate and point (x, r) = (0, 0). The numeric computer analysis of this problem is linked below.

NUMERICS: NB#1

Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (supercritical), pitchfork bifurcation (subcritical).

File content: A quantitatively precise behaviour of the dynamics shown in Fig. 9; the phase portrait and bifurcation diagram corresponding to Eq. (21). Figure 10 shows the resulting diagram.



Figure 10: Bifurcation diagram of the problem given by (21). The stable branch is shown with the solid bold curve and the unstable branch is shown with the dashed curve.

6.1.3 Normal form

Let's take a closer look at the previous result. The series expansion of Eq. (21) is in the form

$$\dot{x} = r + x - \ln(1+x) \approx \begin{bmatrix} \text{Maclaurin ser.} \\ \text{expansion of} \\ \ln(1+x) \\ \text{about } x^* = 0 \end{bmatrix} \approx r + x - \underbrace{\left(x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots\right)}_{\text{the Maclaurin expansion}} \approx r + \frac{x^2}{2} + O(x^3).$$
(29)

This result is similar to the example given by Eq. (17). Surely, after neglecting $O(x^3)$ terms, the qualitative dynamics of this system will be *exactly* the same. Many systems can be reduced to the form (17) or (19) near their respective fixed points. The **normal form** of the saddle-node bifurcation is thus given by

$$\dot{x} = r \pm x^2. \tag{30}$$

All systems that can be reduced to this form share the abovementioned properties.

6.2 Transcritical bifurcation

The normal form of transcritical bifurcation is given by

$$\boxed{\dot{x} = rx \pm x^2,}\tag{31}$$

where r is the control parameter. Let's study the case

$$\dot{x} = rx - x^2 = x(r - x).$$
 (32)

Fixed points for this system are

$$\begin{cases} x_1^* = 0, \quad \forall r, \\ x_2^* = r. \end{cases}$$

$$\tag{33}$$

Figures 11 and 12 show the phase portraits for the varied values of r and the corresponding bifurcation diagram, respectively. Note: $\underline{x^* = 0 \text{ can't be destroyed but its stability can be changed}}$. From Fig. 12 it is clear that the bifurcation coordinate and point (x, r) = (0, 0).



Figure 11: Phase portraits of Eq. (32) for different values of r.



Figure 12: Bifurcation diagram of Eq. (32). The stable branches are shown with the bold continuous lines and the unstable branches are shown with the dashed bold lines.

These result can be compared against the result found using a computer.

NUMERICS: NB#1

Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (supercritical), pitchfork bifurcation (subcritical).

Skip if needed: start

Let's analyse algebraically the stability of the bifurcation diagram branches shown in Fig. 12. In order to determine the stability of the fixed points one needs to analyse the slopes of phase portrait curves at said fixed points x^* :

$$f'(x) = \frac{d}{dx}(rx - x^2) = r - 2x.$$
 (34)

In case of $x_1^* = 0$

$$f'(x_1^*) = f'(0) = r, \quad x_1^* \text{ is } \begin{cases} \text{stable for } r < 0, \\ \text{unstable for } r > 0. \end{cases}$$
(35)

In case of $x_2^* = r$

$$f'(x_2^*) = f'(r) = -r, \quad x_2^* \text{ is } \begin{cases} \text{stable for } r > 0, \\ \text{unstable for } r < 0. \end{cases}$$
(36)



Figure 13: Phase portrait of Eq. (38) for different values of r. In the case r = 0 solutions near the fixed point $x^* = 0$ decay algebraically.



Figure 14: Bifurcation diagram. The stable branches are shown with the bold continuous curves and a lines, and the unstable branch is shown with bold dashed line.

6.3 Pitchfork bifurcation

Pitchfork bifurcations occurs in systems with symmetry. It involves mergers or splitting apart of fixed points and changes in their stabilities. Normal form is given by

$$\dot{x} = rx \pm x^3, \tag{37}$$

where r is the control parameter.

Nonlinear Dynamics

6.3.1 Supercritical pitchfork bifurcation

Let's study the case

$$\dot{x} = rx - x^3. \tag{38}$$

Fixed points for this system are

$$\begin{cases} x_1^* = 0, & \forall r, \\ x_2^* = \pm \sqrt{r}, & r \ge 0. \end{cases}$$
(39)

Figures 13 and 14 show the phase portraits for varied values of r and the corresponding bifurcation diagram, respectively. From Fig. 14 it is clear that the bifurcation coordinate and point (x, r) = (0, 0). The above result can be compared against the result found using a computer.

NUMERICS: NB#1 Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (supercritical), pitchfork bifurcation (subcritical).

6.3.2 Subcritical pitchfork bifurcation

Let's consider the other case of normal form (37) where

$$\dot{x} = rx + x^3. \tag{40}$$

Fixed points for this system are the same as for the previous case but with different stability types. The fixed point are

$$\begin{cases} x_1^* = 0, \quad \forall r, \\ x_2^* = \pm i\sqrt{r}, \quad r < 0, \end{cases}$$

$$\tag{41}$$

where i is the imaginary unit.

Figures 15 and 16 show the phase portraits for varied values of r and the corresponding bifurcation diagram, respectively. From Fig. 16 it is clear that the bifurcation coordinate and point (x, r) = (0, 0).



Figure 15: Phase portrait of Eq. (40) for different values of r. In the case r = 0 the solutions near the fixed point $x^* = 0$ decay algebraically.



Figure 16: Bifurcation diagram. The stable branch is shown with the continuous bold line, and the unstable branches are shown with the dashed bold curves and a line.

The obtained result can be compared against the result found using a computer.

NUMERICS: NB#1

Bifurcation diagrams and 1-D systems: saddle-node bifurcation, transcritical bifurcation, pitchfork bifurcation (subcritical), pitchfork bifurcation (subcritical).

Revision questions

- 1. What does linearisation of a nonlinear system imply?
- 2. Linearise the following 1-D system

$$\dot{x} = x^3 - x \tag{42}$$

- 3. What is bifurcation?
- 4. What is bifurcation diagram?
- 5. What is saddle-node bifurcation?
- 6. What is transcritical bifurcation?
- 7. What is pitchfork bifurcation?
- 8. What is supercritical pitchfork bifurcation?
- 9. What is subcritical pitchfork bifurcation?
- 10. What is normal form in the context of bifurcations?
- 11. Are oscillation possible in 1-D systems?
- 12. Why are oscillations impossible in 1-D systems?
- 13. What does uniqueness of solutions imply in the context of phase space trajectories?